



Normandie Université



UNIVERSITY OF AMSTERDAM

Tobacco smoking and binge drinking among university students in three European countries, 2009-2017

Tavolacci MP1, De Bruyn S2, Van der Heijde C3, Vonk P3, Ladner J4, Van Hal G5.1 CIC 1404, and INSERM 1073, Rouen University Hospital, France. 2Dpt of Sociology, university of Antwerp, Belgium; 3Student Health Service, university of Amsterdam, Netherlands; 4Dpt Of Epidemiology and Health Promotion, Rouen University Hospital, France; 5Social Epidemiology and Health Policy, University of Antwerp, Belgium

OBJECTIVE

Identify the trend of tobacco smoking and binge drinking among university students of Belgium, France and the Netherlands over a 8 years period




3 repeated cross-sectional studies among university students (18-25 years) of Amsterdam (the Netherlands), Antwerp (Belgium), and Rouen (France)

2009-10: 4432 students

2012-13: 6743 students

2016-17: 7515 students

Prevalence of tobacco smoking in the 3 University between 2009 and 2017



63.4% in Antwerp




68.7% in Amsterdam



65.4% in Rouen

Prevalence of binge drinking (>1 / month) in the 3 University between 2009 and 2017



CONCLUSION

Risk behaviours as tobacco smoking and binge drinking in university student show a different prevalence according to country. Tobacco smoking decreases in country with high prevalence, Frequent binge drinking did not change during the 8 years period. Further analyses need to be conducted taking into account the curriculum and the socio-demographic characteristics.